

A. 原著論文

- 1) "Unusually Ready Cleavage of an S-C α Bond in Sterically Crowded Sulphoxides," R. Okazaki, T. Ishida, and N. Inamoto, *J. Chem. Soc., Chem. Commun.*, **1988**, 40-41.
- 2) "Thioaldehyde Anion Radicals," D. Casarini, L. Lunazzi, G. Placucci, T. Ishida, A. Ishii, and R. Okazaki, *J. Org. Chem.*, **53**, 1582-1584 (1988).
- 3) "Bis[3-*tert*-butyl-5-(*N*-oxy-*tert*-butylamino)phenyl] Nitroxide in a Quartet Ground State: A Prototype for Persistent High-Spin Poly[(oxyimino)-1,3-phenylenes]," T. Ishida and H. Iwamura, *J. Am. Chem. Soc.*, **113**, 4238-4241 (1991).
- 4) "An Improved Synthesis of Poly(*m*-aniline) and the Magnetic Properties of Its HCl Salt," T. Ishida and H. Iwamura, *Chem. Lett.*, **1991**, 317-318.
- 5) "Syntheses and Magnetic Properties of Poly(phenylacetylenes) Carrying a (1-Oxido-3-oxy-4,4,5,5-tetramethyl-2-imidazolin-2-yl) Group at the Meta or Para Position of the Phenyl Ring," A. Fujii, T. Ishida, N. Koga, and H. Iwamura, *Macromolecules*, **24**, 1077-1082 (1991).
- 6) "Addition Reaction of Benzyne to C₆₀," M. Tsuda, T. Ishida, T. Nogami, S. Kurono, and M. Ohashi, *Chem. Lett.*, **1992**, 2333-2334.
- 7) "Intramolecular Magnetic Coupling between Two Nitrene or Two Nitroxide Units through 1,1-Diphenylethylene Chromophores. Isomeric Dinitrenes and Dinitroxides Related in Connectivity to Trimethylenemethane, Tetramethyleneethane, and Pentamethylenepropene," T. Matsumoto, T. Ishida, N. Koga, and H. Iwamura, *J. Am. Chem. Soc.*, **114**, 9952-9959 (1992).
- 8) "Intermolecular Ferromagnetic Interaction of 4-(2-Naphthylmethyleneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," K. Tomioka, T. Ishida, T. Nogami, and H. Iwamura, *Chem. Lett.*, **1993**, 625-626.
- 9) "Intermolecular Ferromagnetic Interaction of 4-(Benzylideneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," K. Tomioka, S.-i. Mitsubori, T. Ishida, T. Nogami, and H. Iwamura, *Chem. Lett.*, **1993**, 1239-1242.
- 10) "Isolation and Characterization of Diels-Alder Adducts of C₆₀ with Anthracene and Cyclopentadiene," M. Tsuda, T. Ishida, T. Nogami, S. Kurono, and M. Ohashi, *J. Chem. Soc., Chem. Commun.*, **1993**, 1296-1298.
- 11) "Addition Reactions of Benzyne, Dienes, Dichlorocarbene, and Oxygen to C₆₀," M. Tsuda, T. Ishida, T. Nogami, S. Kurono, and M. Ohashi, *Fullerene Sci. Technol.*, **1**, 275-285 (1993).
- 12) "C₆₁Cl₂. Synthesis and Characterization of Dichlorocarbene Adducts of C₆₀," M. Tsuda, T. Ishida, T. Nogami, S. Kurono, and M. Ohashi, *Tetrahedron Lett.*, **34**, 6911-6912 (1993).
- 13) "A New Organic Ferromagnet: 4-(Benzylideneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," T. Nogami, K. Tomioka, T. Ishida, H. Yoshikawa, M. Yasui, F. Iwasaki, H. Iwamura, N. Takeda, and M. Ishikawa, *Chem. Lett.*, **1994**, 29-32; *Technical Report of ISSP*, Ser. A, No. 2734, Sept. 1993.
- 14) "Organic Second Order Nonlinear Optical Materials with Transparency in Blue Wavelength Region: Ethylene Derivatives Possessing Electron Donor and Acceptor Groups," H. Osaka, T. Ishida, T. Nogami, R. Yamazaki, M. Yasui, F. Iwasaki, A. Mizoguchi, M. Kubata, T. Uemiyama, and A. Nishimura, *Bull. Chem. Soc. Jpn.*, **67**, 918-923 (1994).
- 15) "Pyrimidines as Ferromagnetic Exchange Couplers in Dinuclear Oxovanadium(IV) Complexes," S.-i. Mitsubori, T. Ishida, T. Nogami, and H. Iwamura, *Chem. Lett.*, **1994**, 285-288.
- 16) "Spectral Characterization of Anthracene-, Cyclopentadiene-, and Dichlorocarbene-C₆₀ Adducts," T. Ishida, M. Tsuda, T. Nogami, S. Kurono, and M. Ohashi, *Fullerene Sci. Technol.*, **2**, 155-164 (1994).
- 17) "Fullerene Aziridine. Facile Synthesis and Spectral Characterization of Fullerene Urethane, C₆₀NCO₂CH₂CH₃," T. Ishida, K. Tanaka, and T. Nogami, *Chem. Lett.*, **1994**, 561-562.
- 18) "Synthesis and Structure of Nitrene-C₆₀ Adduct, C₆₀NPhth (Phth = phthalimido)," S.-y. Kuwashima, M. Kubota, K. Kushida, T. Ishida, M. Ohashi, and T. Nogami, *Tetrahedron Lett.*, **35**, 4371-4374 (1994).
- 19) "Molecular Ferromagnet, PM[VO(hfac)₂]₂ (PM = pyrimidine, hfac = 1,1,1,5,5,5-hexafluoropentane-2,4-dionate). Ferromagnetic Assembly of Ground High-Spin Molecules," S.-i. Mitsubori, T. Ishida, T. Nogami, H. Iwamura, N. Takeda, and M. Ishikawa, *Chem. Lett.*, **1994**, 685-686.

- 20) "An Organic Ferromagnet with a T_C of 0.4 K: 4-(*p*-Pheynylbenzylideneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," T. Ishida, H. Tsuboi, T. Nogami, H. Yoshikawa, M. Yasui, F. Iwasaki, H. Iwamura, N. Takeda, and M. Ishikawa, *Chem. Lett.*, **1994**, 919-922;
"An Organic Ferromagnet with a T_C of 0.24 K: 4-(*p*-Pheynylbenzylideneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," T. Nogami, H. Tsuboi, T. Ishida, H. Iwamura, N. Takeda, and M. Ishikawa, *Technical Report of ISSP*, Ser. A, No. 2744, Oct. 1993.
- 21) "Synthesis and Spectral Characterization of Difunctional Fullerenes," T. Ishida, K. Tanaka, T. Furudate, T. Nogami, M. Kubota, S. Kurono, and M. Ohashi, *Fullerene Sci. Technol.*, **3**, 85-92 (1995).
- 22) "Synthesis and Mass Spectral Analysis of C_{60} -Dihalocarbene Adducts," T. Ishida, T. Furudate, T. Nogami, M. Kubota, T. Hirano, and M. Ohashi, *Fullerene Sci. Technol.*, **3**, 399-409 (1995).
- 23) "Syntheses and Structures of Novel Fused-Ring Systems Containing Six or Seven Sulfur Atoms: 2,3:5,6-Bis(ethylenedithio)thieno[3,2-*b*]thiophene and 3,4:6,7-Bis(ethylenedithio)thieno[3,2-*c*][1,2]dithiin," M. Tanaka, T. Ishida, T. Nogami, H. Yoshikawa, M. Yasui, and F. Iwasaki, *Bull. Chem. Soc. Jpn.*, **68**, 1193-1199 (1995).
- 24) "First X-Ray Structural Determination of Fullerene [2+2] Cycloadduct," T. Ishida, K. Shinozuka, T. Nogami, S. Sasaki, and M. Iyoda, *Chem. Lett.*, **1995**, 317-318.
- 25) "Ferromagnetism in Organic Radical Crystal of 4-(*p*-Chlorobenzylideneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," T. Nogami, T. Ishida, H. Tsuboi, H. Yoshikawa, H. Yamamoto, M. Yasui, F. Iwasaki, H. Iwamura, N. Takeda, and M. Ishikawa, *Chem. Lett.*, **1995**, 635-636; *Technical Report of ISSP*, Ser. A, No. 2932, Feb. 1995.
- 26) "Fullerene Spin Label. Synthesis and Characterization of the [60]Fullerene-substituted TEMPO Radical," T. Ishida, K. Shinozuka, M. Kubota, M. Ohashi, and T. Nogami, *J. Chem. Soc., Chem. Commun.*, **1995**, 1841-1842.
- 27) "An Organic Metamagnet, 4-(2-Naphthylmethyleneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," T. Ishida, K. Tomioka, T. Nogami, H. Yoshikawa, M. Yasui, F. Iwasaki, H. Iwamura, N. Takeda, and M. Ishikawa, *Chem. Phys. Lett.*, **247**, 7-12 (1995).
- 28) "Synthesis and Characterization of C_{60} Derivatives Possessing TEMPO Radicals," T. Ishida, K. Shinozuka, T. Nogami, M. Kubota, and M. Ohashi, Symposium-in-Print on "Fullerene Chemistry," *Tetrahedron*, **52**, 5103-5112 (1996).
- 29) "Synthesis and Characterization of Dialkyl- C_{61} ," M. Kawasumi, T. Ishida, and T. Nogami, *Fullerene Sci. Technol.*, **4**, 357-367 (1996).
- 30) "Synthesis, Structure, and Some Reactions of (2,4,6-tri-*t*-Butyl)thiobenzaldehyde, the First Stable Aromatic Thioaldehyde," A. Ishii, T. Ishida, N. Kumon, N. Fukuda, H. Oyama, N. Inamoto, F. Iwasaki, and R. Okazaki, *Bull. Chem. Soc. Jpn.*, **69**, 709-717 (1996).
- 31) "Proposed Mechanism of Ferromagnetic Interaction of Organic Ferromagnets: 4-(Arylmethyleneamino)-2,2,6,6-tetramethylpiperidin-1-oxyls and Related Compounds," T. Nogami, T. Ishida, M. Yasui, F. Iwasaki, N. Takeda, M. Ishikawa, T. Kawakami, and K. Yamaguchi, *Bull. Chem. Soc. Jpn.*, **69**, 1841-1848 (1996).
- 32) "Unusual Reactions of C_{60} with Aldehydes in the Presence of Aqueous Ammonia," A. Komori, M. Kubota, T. Ishida, H. Niwa, M. Ohashi, and T. Nogami, *Tetrahedron Lett.*, **37**, 4031-4034 (1996).
- 33) "Organic Radicals Exhibiting Intermolecular Ferromagnetic Interactions with High Probability: 4-Arylmethyleneamino-2,2,6,6-tetramethylpiperidin-1-yloxyyls and Related Compounds," K. Togashi, R. Imachi, K. Tomioka, H. Tsuboi, T. Ishida, T. Nogami, N. Takeda, and M. Ishikawa, *Bull. Chem. Soc. Jpn.*, **69**, 2821-2830 (1996).
- 34) "TEMPO-Based Organic Ferromagnets and Metamagnets (TEMPO = 2,2,6,6-Tetramethylpiperidin-1-yloxyyl)," T. Nogami and T. Ishida, *Mol. Cryst. Liq. Cryst.*, **296**, 305-322 (1997).
- 35) " μ SR of the Organic Radical Ferromagnet, 4-(*p*-Chlorobenzylideneamino)-2,2,6,6-tetramethylpiperidin-1-yloxyyl," R. Imachi, T. Ishida, T. Nogami, S. Ohira, K. Nishiyama, and K. Nagamine, *Chem. Lett.*, **1997**, 233-234.

- 36) "Ferromagnetic Dimer of π -Conjugated Organic Free Radicals 9,9-Bis(*p*-tolyl)-9,10-dihydroacridin-10-yloxy," R. Imachi, T. Ishida, M. Suzuki, M. Yasui, F. Iwasaki, and T. Nogami, *Chem. Lett.*, **1997**, 743-744.
- 37) "Pyrimidine-Bridged Cobalt(II) Complexes with a Chiral 3-D Network Showing Weak Ferromagnetism," K. Nakayama, T. Ishida, R. Takayama, D. Hashizume, M. Yasui, F. Iwasaki, and T. Nogami, *Chem. Lett.*, **1998**, 497-498.
- 38) "Electrical Conductivities and Crystal and Band-Electronic Structures of a New Phase of BEDT-TTF-Bromide Salt, (BEDT-TTF)₂Br(H₂O)₃," M.-Y. Luo, T. Ishida, A. Kobayashi, and T. Nogami, *Synth. Met.*, **96**, 97-102 (1998).
- 39) "Ferromagnetic and Antiferromagnetic Phases of 4-(4-Iodobenzylideneamino)-2,2,6,6-tetramethylpiperidin-1-yloxy," T. Ishida, K. Togashi, R. Imachi, H. Yamamoto, M. Yasui, F. Iwasaki, and T. Nogami, *Bull. Chem. Soc. Jpn.*, **71**, 2583 (1998).
- 40) "Solid-State High-Resolution ¹H and ²D NMR Study of the Electron Spin Distribution of the Hydrogen-Bonded Organic Ferromagnetic Compound 4-Hydroxyimino-TEMPO," G. Maruta, S. Takeda, R. Imachi, T. Ishida, T. Nogami, and K. Yamaguchi, *J. Am. Chem. Soc.*, **121**, 424-431 (1999).
- 41) "Intermolecular Ferromagnetic Interaction in the Crystal of a Diphenyl Nitroxide Derivative. The Role of Spin-Polarized Hydrogen Atoms Located near a Neighboring N-O Site," M. Nakagawa, T. Ishida, M. Suzuki, D. Hashizume, M. Yasui, F. Iwasaki, and T. Nogami, *Chem. Phys. Lett.*, **302**, 125-131 (1999).
- 42) "TEMPO Radicals Showing Magnetic Interactions. I. 4-(4-Halobenzylideneamino)-TEMPO and Related Compounds," F. Iwasaki, J. H. Yoshikawa, H. Yamamoto, E. Kan-nari, K. Takada, M. Yasui, T. Ishida, and T. Nogami, *Acta Crystallogr., Sect. B*, **55**, 231-245 (1999).
- 43) "TEMPO Radicals Showing Magnetic Interactions. II. 4-(Benzylideneamino)-TEMPO and Related Compounds," F. Iwasaki, J. H. Yoshikawa, H. Yamamoto, K. Takada, E. Kan-nari, M. Yasui, T. Ishida, and T. Nogami, *Acta Crystallogr., Sect. B*, **55**, 1057-1067 (1999).
- 44) "Dichlorobis(pyrimidine-*N*)cobalt(II) and Its Bromo Derivative," D. Hashizume, R. Takayama, K. Nakayama, T. Ishida, T. Nogami, M. Yasui, and F. Iwasaki, *Acta Crystallogr., Sect. C*, **55**, 1793-1797 (1999).
- 45) "Molecular Orbital Study on the Magnetic Properties of Pyrazine- and Pyrimidine-Bridged Copper(II) Complexes," F. Mohri, K. Yoshizawa, T. Yamabe, T. Ishida, and T. Nogami, *Molecular Engineering*, **8**, 357-373 (1999).
- 46) "Low-Temperature Heat Capacities and Ferromagnetic Phase Transition of the Organic Free Radical Ferromagnet, 4-(4-Chlorobenzylideneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl (CATMP)," Y. Miyazaki, T. Matsumoto, T. Ishida, T. Nogami, and M. Sorai, *Bull. Chem. Soc. Jpn.*, **73**, 67-71 (2000).
- 47) "First TOF Donors. Synthesis and Characterization of Dibenzo- and Dinaphthotetraoxafulvalenes," K. Tanaka, K. Yoshida, T. Ishida, A. Kobayashi, and T. Nogami, *Adv. Mater.*, **12**, 661-664 (2000).
- 48) "The Experimental Deformation Electron-Density Distributions of 4-Arylmethyleneamino-2,2,6,6-tetramethyl-1-piperidyloxy (TEMPO) Radicals: Relations between Electron Densities and Intermolecular Interactions," M. Yasui, E. Kan-nari, T. Ishida, T. Nogami, and F. Iwasaki, *Bull. Chem. Soc. Jpn.*, **73**, 1333-1340 (2000).
- 49) "Spin Density of a Ferromagnetic Tempo Derivative: Polarized Neutron Investigation and Ab Initio Calculation," Y. Pontillon, A. Grand, T. Ishida, E. L. Berna, T. Nogami, E. Ressouche, and J. Schweizer, *J. Mater. Chem.*, **10**, 1539-1546 (2000).
- 50) "One-Dimensional $S = 1$ Ferromagnetic Chain Formed by a Square Ni₂S₂ Motif in Ni(qt)₂ (qt = Quinoline-8-thiolate)," T. Miyake, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Chem. Lett.*, **2000**, 952-953.
- 51) "Low-Temperature Magnets M[N(CN)₂]₂(pyrimidine) (M = Fe and Co) with a 3-D Network," T. Kusaka, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Chem. Lett.*, **2000**, 1146-1147.
- 52) "Zero-Field Muon Spin Rotation Study on Genuine Organic Ferromagnets, 4-Arylmethyleneamino-2,2,6,6-tetramethylpiperidin-1-yloxy (Ar = 4-biphenyl and phenyl)," T. Ishida, S. Ohira, T. Ise, K. Nakayama, I. Watanabe, T. Nogami, and K. Nagamine, *Chem. Phys. Lett.*, **330**, 110-117 (2000).

- 53) "Magnetic-Field Induced Gap and Staggered Susceptibility in the $S = 1/2$ Chain $[\text{PM}^+\text{Cu}(\text{NO}_3)_2(\text{H}_2\text{O})_2]_n$ (PM = Pyrimidine)," R. Feyerherm, S. Abens, D. Günther, T. Ishida, M. Meissner, M. Meschke, T. Nogami, and M. Steiner, *J. Phys.: Condens. Matter*, **12**, 8495-8509 (2000).
- 54) "Pressure Effects of Genuine Organic Crystalline Ferromagnet Possessing Intermolecular Contacts between Nitroxide Oxygen and Methyl Hydrogen Atoms," M. Mito, T. Kawae, M. Hitaka, K. Takeda, T. Ishida, and T. Nogami, *Chem. Phys. Lett.*, **333**, 69-75 (2001).
- 55) "Single-Component Organic Semiconductors Based on Novel Radicals that Exhibit Electrochemical Amphotericity: Preparation, Crystal Structures, and Solid-State Properties of N,N' -Dicyanopyrazinonaphthoquinodiiminides Substituted with an N -Alkylpyrimidinium Unit," T. Suzuki, S. Miyanari, Y. Tsubata, T. Fukushima, T. Miyashi, Y. Yamashita, K. Imaeda, T. Ishida, and T. Nogami, *J. Org. Chem.*, **66**, 216-224 (2001).
- 56) "Dipyrimidine-Cu(II) Dinitrate Complexes Showing Magnetic Interactions," M. Yasui, Y. Ishikawa, N. Akiyama, T. Ishida, T. Nogami, and F. Iwasaki, *Acta Crystallogr., Sect. B*, **57**, 288-295 (2001).
- 57) "Radical-Copper Wheels: Structure and Magnetism of Hexanuclear Hybrid Arrays," J. Omata, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Inorg. Chem.*, **40**, 3954-3958 (2001).
- 58) "Ferromagnetic Exchange Coupling of Vanadium(IV) $d\pi$ Spins across Pyrimidine Rings: Dinuclear Complexes of Oxovanadium(IV) Bis(1,1,1,5,5,5-hexafluoropentane-2,4-dionate) Bridged by Pyrimidine Derivatives," T. Ishida, S. -i. Mitsubori, T. Nogami, N. Takeda, M. Ishikawa, and H. Iwamura, *Inorg. Chem.*, **40**, 7059-7064 (2001).
- 59) "Bis(hfac)-Copper(II) Complexes Bridged by Pyrimidines Showing Magnetic Interactions," M. Yasui, Y. Ishikawa, T. Ishida, T. Nogami, and F. Iwasaki, *Acta Crystallogr., Sect. B*, **57**, 772-779 (2001).
- 60) "Novel Organic-Inorganic Molecular Assembly: a Copper(I) and Copper(II) Mixed-Valent Cluster with Bromide Bridges and Paramagnetic Ligands," T. Katayama, T. Ishida, and T. Nogami, *Inorg. Chim. Acta*, **329**, 31-35 (2002).
- 61) "Low-Temperature Heat Capacity and One-Dimensional Ferromagnetic Behavior of the Organic Free Radical, 4-Benzylideneamino-2,2,6,6-tetramethylpiperidin-1-oxyl (BATMP)," T. Sakakibara, Y. Miyazaki, T. Ishida, T. Nogami, and M. Sorai, *J. Phys. Chem. B*, **106**, 6390-6394 (2002).
- 62) "Antiferromagnetic Coupling of Transition Metal Spins across Pyrimidine and Pyrazine Bridges in Dinuclear Manganese(II), Cobalt(II), Nickel(II), and Copper(II) 1,1,1,5,5,5-Hexafluoropentane-2,4-dionate Complexes," T. Ishida, T. Kawakami, S.-i. Mitsubori, T. Nogami, K. Yamaguchi, and H. Iwamura, *J. Chem. Soc., Dalton Trans.*, **2002**, 3177-3186.
- 63) "Convenient Synthesis and Host-Guest Compounds of 9,9'(10*H*,10'*H*)-Spirobiacridines," M. Ooishi, M. Seino, R. Imachi, T. Ishida, and T. Nogami, *Tetrahedron Lett.*, **43**, 5521-5524 (2002).
- 64) "Metal-Azide-Pyrimidine Complexes $\text{M}(\text{N}_3)_2(\text{pm})$ with a Three-Dimensional Network Showing Weak Ferromagnetism for $\text{M} = \text{Mn}$ and Fe and Antiferromagnetism for $\text{M} = \text{Co}$ and Ni ," Y. Doi, T. Ishida, and T. Nogami, *Bull. Chem. Soc. Jpn.*, **75**, 2455-2461 (2002).
- 65) "Crystal Structures and Magnetic Properties of Copper(II) Hexafluoroacetylacetonate Complexes with hnn and hin and Manganese(II) Hexafluoroacetylacetonate Complexes with hin (hnn = 4,4,5,5-Tetramethylimidazolin-1-oxyl 3-Oxide; hin = 4,4,5,5-Tetramethylimidazolin-1-oxyl)," T. Ise, T. Ishida, and T. Nogami, *Bull. Chem. Soc. Jpn.*, **75**, 2463-2468 (2002).
- 66) "Nitronyl Nitroxide Substituted Aniline *o*-APNN with a Three-Dimensional Hydrogen Bond Network Showing Ferromagnetic Interaction," K. Doi, T. Ishida, and T. Nogami, *Chem. Lett.*, **32**, 544-545 (2003).
- 67) "Metamagnetic Behavior of $[\text{Ni}(\text{4ImNNH})_2(\text{NO}_3)_2]$ Having a Ground High-Spin State (4ImNNH = 4-Imidazolyl Nitronyl Nitroxide)," C. Aoki, T. Ishida, and T. Nogami, *Inorg. Chem. Commun.*, **6**, 1122-1125 (2003).
- 68) "Rectangular Tetragadolium Complex Ion with a Ground $S = 27/2$ State Containing Bis(dimethylglyoximate)copper(II) as a Ferrimagnetic Exchange Coupler," Y. Kobayashi, S. Ueki, T. Ishida, and T. Nogami, *Chem. Phys. Lett.*, **378**, 337-342 (2003).
- 69) "Ferro- and Ferrimagnetic Chains of hin-Bridged Copper(II) and Manganese(II) and hnn-Bridged Manganese(II) Complexes (hin = 4,4,5,5-Tetramethylimidazolin-1-oxyl; hnn =

- 4,4,5,5-Tetramethylimidazolin-1-oxyl 3-Oxide)," T. Ise, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Inorg. Chem.*, **42**, 6106-6113 (2003).
- 70) "A Self-Assembled Helix from 4,6-Dimethylpyrimidine and Copper(II) Bromide," T. Ishida, L. Yang, and T. Nogami, *Chem. Lett.*, **32**, 1018-1019 (2003).
- 71) "Molecular Metamagnet $[\text{Ni}(\text{4ImNNH})_2(\text{NO}_3)_2]$ (4ImNNH = 4-Imidazolyl Nitronyl Nitroxide) and the Related Compounds Showing Supramolecular H-Bonding Interaction," C. Aoki, T. Ishida, and T. Nogami, *Inorg. Chem.*, **42**, 7616-7625 (2003).
- 72) "Supramolecular Triangular and Linear Arrays of Metal-Radical Solids Using Imino-Nitroxide-Substituted Pyrazolate Bridges," S. Yamada, T. Ishida, and T. Nogami, *Dalton Trans.*, **2004**, 898-903.
- 73) "Weak Ferromagnetism with Very Large Canting in a Chiral Lattice: $\text{Fe}(\text{pyrimidine})_2\text{Cl}_2$," R. Feyerherm, A. Loose, T. Ishida, T. Nogami, J. Kreitlow, D. Baabe, F. J. Litterst, S. Söllow, H.-H. Klauss, and K. Doll, *Phys. Rev. B*, **69**, 134427(1)-134427(8) (2004).
- 74) "Single-Crystal Magnetic Study on Guest-Tunable Weak Ferromagnets $[\text{M}\{\text{N}(\text{CN})_2\}_2(\text{pyrimidine})](\text{guest})$ (M = Fe, Co)," N. Takagami, T. Ishida, and T. Nogami, *Bull. Chem. Soc. Jpn.*, **77**, 1125-1134 (2004). (Selected Paper)
- 75) "Crystal Structure of a Molecular Complex from Native β -Cyclodextrin and Copper(II) Chloride," G. Kurokawa, M. Sekii, T. Ishida, and T. Nogami, *Supramol. Chem.*, **16**, 381-384 (2004).
- 76) "Remarkably Strong Intermolecular Antiferromagnetic Couplings in the Crystal of Biphenyl-3,5-diyl Bis(*tert*-butyl Nitroxide)," G. Kurokawa, T. Ishida, and T. Nogami, *Chem. Phys. Lett.*, **392**, 74-79 (2004).
- 77) "Dinuclear Manganese(II) Complex Containing Tetrakis(2-pyridyl)methane as a Spiro-fused Bridge," A. Okazawa, T. Ishida, and T. Nogami, *Chem. Lett.*, **33**, 1478-1479 (2004).
- 78) "Magnetic Coupling Switch by Means of Host-Guest Chemistry in a Polyether Side-Arm," T. Nakabayashi, T. Ishida, and T. Nogami, *Inorg. Chem. Commun.*, **7**, 1221-1224 (2004).
- 79) "Synthesis and Characterization of Dibenzodioxadiselenafulvalene," T. Kojima, K. Tanaka, T. Ishida, and T. Nogami, *J. Org. Chem.*, **69**, 9319-9322 (2004).
- 80) "Structure and Magnetic Properties of a Honeycomb-like Copper(II) Complex Containing 4-Aminopyrimidine and Methoxide Bridges," K. Osanai, T. Ishida, and T. Nogami, *Chem. Lett.*, **34**, 620-621 (2005).
- 81) "Pressure Effect on the Magnetic Phase Transition of Guest-Tunable Weak Ferromagnets $[\text{Fe}\{\text{N}(\text{CN})_2\}_2(\text{pyrimidine})](\text{guest})$ (Guest = Ethanol and Pyrimidine)," K. Yuza, T. Ishida, and T. Nogami, *Chem. Lett.*, **34**, 974-975 (2005).
- 82) "Trans-fused Macrobicyclic "Betweenanene" around a Planar $\text{Cu}_2(\mu\text{-CH}_3\text{O})_2$ Core Showing Remarkable Antiferromagnetic Interaction," T. Ishida, K. Yanagi, and T. Nogami, *Inorg. Chem.*, **44**, 7307-7309 (2005).
- 83) "The First A_2B_2 -Heterometal Ferrimagnetic Chain. Structures and Magnetic Properties of Polymeric $[\text{Gd}_2\text{Cu}_2]_n$ and the Corresponding Monomer," S. Ueki, Y. Kobayashi, T. Ishida, and T. Nogami, *Chem. Commun.*, **2005**, 5223-5225.
- 84) "Organic Kagomé Lattice Consisting of Trimeric Pyrazole and Oxime Supramolecular Synthons from 3-Pyrazolecarboxaldoxime," T. Nyui, T. Nogami, and T. Ishida, *CrystEngComm*, **7**, 612-615 (2005).
- 85) "Self-assembled *meso*-Helicates of Linear Trinuclear Nickel(II)-Radical Complexes with Triple Pyrazolate Bridges," S. Yamada, M. Yasui, T. Nogami, and T. Ishida, *Dalton Trans.*, 1622-1626 (2006).
- 86) "Bio-Inspired Materials for Electron Donors and Charge-Transfer Complex Formation Based on the Imidazopyrazinone Luciferin Core," M. Suzuki, T. Fujii, T. Nogami, T. Hirano, and T. Ishida, *Chem. Lett.*, **35**, 250-251 (2006).
- 87) "Oximate-Bridged Trinuclear Dy-Cu-Dy Complex Behaving as a Single-Molecule Magnet and Its Mechanistic Investigation," F. Mori, T. Nyui, T. Ishida, T. Nogami, K.-Y. Choi, and H. Nojiri, *J. Am. Chem. Soc.*, **128**, 1440-1441 (2006).
- 88) "Polymorphic Alternating HNN-Cobalt(II) Chains Both Behaving as Single-Chain Magnets (HNN = 4,4,5,5-Tetramethylimidazolin-1-oxyl 3-Oxide)," N. Ishii, T. Ishida, and T. Nogami, *Inorg. Chem.*, **45**, 3837-3839 (2006).
- 89) "Illumination-induced Magnetization Enhancement of the Weak Ferromagnet $[\text{Fe}(\text{N}_3)_2(\text{pm})]$ (pm = Pyrimidine)," Y. Morishita, Y. Doi, T. Nogami, and T. Ishida, *Chem. Lett.*, **35**, 770-771 (2006).

- 90) "Bistable Polymorphs Showing Diamagnetic and Paramagnetic States of an Organic Crystalline Biradical Biphenyl-3,5-diyl Bis(*tert*-butylnitroxide)," H. Nishimaki, S. Mashiyama, M. Yasui, T. Nogami, and T. Ishida, *Chem. Mater.*, **18**, 3602-3604 (2006).
- 91) "Ferromagnetic Exchange Coupling in a Spiro-Fused Dinuclear Nickel(II) Complex with an Approximate D_{2d} Symmetry," A. Okazawa, T. Nogami, and T. Ishida, *Chem. Phys. Lett.*, **427**, 333-337 (2006).
- 92) "Double Half-cubane Copper(II) Complexes Available from Copper(II) 1,1,1,5,5,5-Hexafluoropentane-2,4-dionate Dissolved in Formamides," Y. Morishita, T. Kogane, T. Nogami, and T. Ishida, *Dalton Trans.*, 4438-4443 (2006).
- 93) "Strong Ferromagnetic Exchange Couplings in Copper(II) and Nickel(II) Complexes with a Paramagnetic Tridentate Chelate Ligand, 2,2'-Bipyridin-6-yl *tert*-Butyl Nitroxide," K. Osanai, A. Okazawa, T. Nogami, and T. Ishida, *J. Am. Chem. Soc.*, **128**, 14008-14009 (2006).
- 94) "Magnetic Exchange Coupling Tunable by Means of Selective Cation Binding into Poly(radical-armed) Azacrowns," K. Igarashi, T. Nogami, and T. Ishida, *Chem. Commun.*, **2007**, 501-503.
- 95) "*tert*-Butyl 2-Pyridyl Nitroxide Available as a Paramagnetic Chelate Ligand for Strongly Exchange-Coupled Metal-Radical Compounds," A. Okazawa, T. Nogami, and T. Ishida, *Chem. Mater.*, **19**, 2733-2735 (2007).
- 96) "Quantum Tunneling of Magnetization via Well-Defined Dy-Cu Exchange Coupling in a Ferrimagnetic High-Spin $[Dy_4Cu]$ Single-Molecule Magnet," S. Ueki, T. Ishida, T. Nogami, K.-Y. Choi, and H. Nojiri, *Chem. Phys. Lett.*, **440**, 263-267 (2007).
- 97) "Giant Coercivity in a One-Dimensional Cobalt-Radical Coordination Magnet," N. Ishii, Y. Okamura, S. Chiba, T. Nogami, and T. Ishida, *J. Am. Chem. Soc.*, **130**, 24-25 (2008).
- 98) "Exchange Coupling and Energy-Level Crossing in a Magnetic Chain $[Dy_2Cu_2]_n$ Evaluated by High-Frequency Electron Paramagnetic Resonance," A. Okazawa, T. Nogami, H. Nojiri, and T. Ishida, *Chem. Mater.*, **20**, 3110-3119 (2008).
- 99) "Pyrazine-Bridged Linear Pentanuclear Copper(II) Complex and Related Tri- and Dinuclear Complexes Showing Various Coordination Structures and Magnetic Couplings," N. Koyama, T. Ishida, T. Nogami, and T. Kogane, *Polyhedron*, **27**, 2341-2348 (2008).
- 100) "Spin-crossover Transition Coupled with Another Solid-Solid Phase Transition for Iron(II) Thiocyanate Complexes Chelated with Alkylated *N*-(Di-2-pyridylmethylene)anilines," Y. Oso, D. Kanatsuki, S. Saito, T. Nogami, and T. Ishida, *Chem. Lett.*, **37**, 760-761 (2008).
- 101) "Magneto-Structure Relationship in Copper(II) and Nickel(II) Complexes Chelated with Stable *tert*-Butyl 5-Phenyl-2-pyridyl Nitroxide and Related Radicals," A. Okazawa, Y. Nagaichi, T. Nogami, and T. Ishida, *Inorg. Chem.*, **47**, 8859-8868 (2008).
- 102) "Ferromagnetic Dy-Ni and Antiferromagnetic Dy-Cu Couplings in Single-Molecule Magnets $[Dy_2Ni]$ and $[Dy_2Cu]$," A. Okazawa, T. Nogami, H. Nojiri, and T. Ishida, *Inorg. Chem.*, **47**, 9763-9765 (2008).
Correction: A. Okazawa, T. Nogami, H. Nojiri, and T. Ishida, *Inorg. Chem.*, **48**, 3292 (2009).
- 103) "Crystal Structures and Magnetic Properties of Two Phases from Imidazolate-Bridged Copper(II) 4,6-Nonanedionate Complexes," R. Watanabe, N. Koyama, T. Nogami, T. Ishida, and T. Kogane, *Chem. Lett.*, **38**, 30-31 (2009).
- 104) "Ferromagnetic Superexchange Coupling through a Low-Spin Iron(II) Ion in a Mixed-Valent Iron(III, II, III) *Meso*-Helicate," T. Mochizuki, T. Nogami, and T. Ishida, *Inorg. Chem.*, **48**, 2254-2259 (2009).
- 105) "Spin-crossover Transition in a *Meso*-phase Iron(II) Thiocyanate Complex Chelated with 4-Hexadecyl-*N*-(2-pyridylmethylene)aniline," Y. Oso and T. Ishida, *Chem. Lett.*, **38**, 604-605 (2009).
- 106) "The Hardest Magnet from a Coordination Polymer of Alternating Cobalt(II)-Radical Heterospins," Y. Okamura, T. Nogami, and T. Ishida, *Chem. Lett.*, **38**, 740-741 (2009).
- 107) "Long-Range Order in the One-Dimensional Cobalt(II)-Radical Coordination Polymer Magnet Probed by Muon Spin Rotation and Relaxation," T. Ishida, Y. Okamura, and I. Watanabe, *Inorg. Chem.*, **48**, 7012-7014 (2009).
- 108) "Super-superexchange Coupling through a Hydrogen Bond in a Linear Copper(II) Complex, $[Cu(LH)(L)]\cdot BF_4\cdot 2H_2O$ (LH = *N-tert*-Butyl-*N*-2-pyridylhydroxylamine)," A. Okazawa and T. Ishida, *Chem. Phys. Lett.*, **480**, 198-202 (2009).

- 109) "Amino Acid Spin-Labels. An Application of Chelation Ability to a Nickel(II) Ion," S. Osada, K. Igarashi, T. Nogami, and T. Ishida, *Chem. Lett.*, **39**, 576-577 (2010).
- 110) "Hard Magnets after Freezing of Spin Dynamics of Soft Magnets in Cobalt(II)-Radical Chain Compounds," Y. Okamura, N. Ishii, T. Nogami, and T. Ishida, *Bull. Chem. Soc. Jpn.*, **83**, 716-725 (2010).
- 111) "Organic Two-Step Spin-Transition-Like Behavior in a Linear $S = 1$ Array: 3'-Methylbiphenyl-3,5-diyl Bis(*tert*-butylnitroxide) and Related Compounds," H. Nishimaki and T. Ishida, *J. Am. Chem. Soc.*, **132**, 9598-9599 (2010).
- 112) "Ferro- and Antiferromagnetic Coupling Switch Accompanied by Twist Deformation around the Copper(II) and Nitroxide Coordination Bond," A. Okazawa, D. Hashizume, and T. Ishida, *J. Am. Chem. Soc.*, **132**, 11516-11524 (2010).
- 113) "Spin-Transition-Like Behavior on One Side in a Nitroxide-Copper(II)-Nitroxide Triad System," A. Okazawa and T. Ishida, *Inorg. Chem.*, **49**, 10144-10147 (2010).
- 114) "Ferromagnetic Gd-Cu, Tb-Cu, and Ho-Cu Couplings in Isomorphous $[Ln_2Cu]$ Complexes," A. Okazawa, R. Watanabe, M. Nezu, T. Shimada, S. Yoshii, H. Nojiri, and T. Ishida, *Chem. Lett.*, **39**, 1331-1332 (2010).
- 115) "Biradical Chelating Host 2,2'-Bipyridine-6,6'-diyl Bis(*tert*-Butyl Nitroxide) Showing Tunable Exchange Magnetic Coupling," K. Koide and T. Ishida, *Inorg. Chem. Commun.*, **14**, 194-196 (2011).
- 116) "Chemical Trend of 4f-3d Exchange Couplings in Heterometallic Complexes with Ln = Gd, Td, Dy, Ho, Er and M = Cu, V," R. Watanabe, K. Fujiwara, A. Okazawa, G. Tanaka, S. Yoshii, H. Nojiri, and T. Ishida, *Chem. Commun.*, **47**, 2110-2112 (2011).
- 117) "Structure and Magnetic Properties of an Unprecedented *syn-anti* μ -Nitrito-1 κ O:2 κ O' Bridged Mn(III)-Salen Complex and Its Isoelectronic and Isostructural Formate Analogue," P. Kar, R. Biswas, M. G. B. Drew, Y. Ida, T. Ishida, and A. Ghosh, *Dalton Trans.*, **40**, 3295-3304 (2011).
- 118) "Spin Canted Antiferromagnetic Phase Transition in Alternating Phenoxo and Carboxylato Bridged Mn(III)-Salen Complexes," P. Kar, P. Mukherjee, M. G. B. Drew, T. Ishida, and A. Ghosh, *Eur. J. Inorg. Chem.*, 2075-2085 (2011).
- 119) "Ferromagnetic Exchange Couplings Showing a Chemical Trend in Cu-Ln-Cu Complexes (Ln = Gd, Tb, Dy, Ho, Er)," T. Shimada, A. Okazawa, N. Kojima, S. Yoshii, H. Nojiri, and T. Ishida, *Inorg. Chem.*, **50**, 10555-10557 (2011).
- 120) "A Unique Example of Structural Diversity Tuned by Apparently Innocent *o*-, *m*- and *p*-Nitro Substituent of Benzoate in Their Complexes of Mn(II) with 4,4'-Bipyridine: 1D Ladder, 2D Sheet and 3D Framework," P. Kar, R. Biswas, Y. Ida, T. Ishida, and A. Ghosh, *Cryst. Growth Design*, **11**, 5305-5315 (2011).
- 121) "Triply Radical-coordinated Gadolinium(III) Complex as a High-spin $S = 5$ Assembly," N. Ikegaya, T. Kanetomo, R. Murakami, and T. Ishida, *Chem. Lett.*, **41**, 82-83 (2012).
- 122) "Oximate-bridged Light-lanthanoid Ln_4Cu Complexes Showing Ferromagnetic Ln-Cu Exchange Coupling," K. Fujiwara, A. Okazawa, G. Tanaka, S. Yoshii, H. Nojiri, and T. Ishida, *Chem. Phys. Lett.*, **530**, 49-54 (2012).
- 123) "Polyether-Bridged Bis(*tert*-Butyl Nitroxide) Paramagnetic Hosts Showing Receptor Ability to Calcium(II) and Barium(II) Ions," S. Osada, N. Hirosawa, and T. Ishida, *Tetrahedron*, **68**, 6193-6197 (2012).
- 124) "Spin-Crossover and Light-Induced Excited Spin-State Trapping Observed for an Iron(II) Complex Chelated with Tripodal Tetrakis(2-pyridyl)methane," N. Hirosawa, Y. Oso, and T. Ishida, *Chem. Lett.*, **41**, 716-718 (2012).
- 125) "Exchange Coupling in TbCu and DyCu Single-molecule Magnets and Related Lanthanide and Vanadium Analogs," T. Ishida, R. Watanabe, K. Fujiwara, A. Okazawa, N. Kojima, G. Tanaka, S. Yoshii, and H. Nojiri, *Dalton Trans.*, **41**, 13609-13619 (2012) (Journal Back Cover).
- 126) "Formation of Two Drastically Different MOFs Based on Mn(II)-Benzoate and Pyrazine with a Change in Seasonal Temperature: Structural Analysis and Magnetic Study," P. Kar, Y. Ida, T. Ishida, and A. Ghosh, *CrystEngComm*, **15**, 400-410 (2013).
- 127) "A New Family of Trinuclear Nickel(II) Complexes as Single-Molecule Magnets," R. Biswas, Y. Ida, M. L. Baker, S. Biswas, P. Kar, H. Nojiri, T. Ishida, and A. Ghosh, *Chem. Eur. J.*, **19**, 3943-3953 (2013).

- 128) “Ground Triplet Pyrimidine-4,6-diyl Bis(*tert*-butyl Nitroxide) as a Paramagnetic Building Block for Metal-Organic Frameworks,” Y. Homma, A. Okazawa, and T. Ishida, *Tetrahedron Lett.*, **54**, 3120-3123 (2013).
- 129) “Syntheses, Structures, and Magnetic Properties of Acetato- and Diphenolato-Bridged 3d-4f Binuclear Complexes $[M(3\text{-MeOsalt})_x(\text{MOH})_x(\text{ac})\text{Ln}(\text{hfac})_2]$ ($M = \text{Zn}^{\text{II}}, \text{Cu}^{\text{II}}, \text{Ni}^{\text{II}}, \text{Co}^{\text{II}}$; $\text{Ln} = \text{La}^{\text{III}}, \text{Gd}^{\text{III}}, \text{Tb}^{\text{III}}, \text{Dy}^{\text{III}}$; 3-MeOsalt = *N,N'*-Bis(3-methoxy-2-oxybenzylidene)-1,3-propanediaminato; ac = Acetato; hfac = Hexafluoroacetylacetonato; $x = 0$ or 1),” M. Towatari, K. Nishi, T. Fujinami, N. Matsumoto, Y. Sunatsuki, M. Kojima, N. Mochida, T. Ishida, N. Re, and J. Mrozinski, *Inorg. Chem.*, **52**, 6160-6178 (2013).
- 130) “A Supramolecular Switch between Ground High- and Low-Spin States Using 2,2':6',2"-Terpyridine-6,6"-Diyl Bis(*tert*-butyl Nitroxide),” T. Konno, K. Koide, and T. Ishida, *Chem. Commun.*, **49**, 5156-5158 (2013).
- 131) “Carbonate-Bridged $\text{Ni}^{\text{II}}_2\text{Ln}^{\text{III}}_2$ ($\text{Ln}^{\text{III}} = \text{Gd}^{\text{III}}, \text{Tb}^{\text{III}}, \text{Dy}^{\text{III}}$) Complexes Generated by Atmospheric CO_2 Fixation and Their Single-Molecule-Magnet Behavior: $[(\mu_4\text{-CO}_3)_2\{\text{Ni}^{\text{II}}(3\text{-MeOsalt})(\text{MeOH}$ or $\text{H}_2\text{O})\text{Ln}^{\text{III}}(\text{NO}_3)_2\}_2]\cdot\text{solvent}$ (3-MeOsalt = *N,N'*-Bis(3-methoxy-2-oxybenzylidene)-1,3-propanediaminato),” S. Sakamoto, T. Fujinami, K. Nishi, N. Matsumoto, N. Mochida, T. Ishida, Y. Sunatsuki, and N. Re, *Inorg. Chem.*, **52**, 7218-7229 (2013).
- 132) “Single-molecule magnet $[\text{Tb}(\text{hfac})_3(2\text{pyNO})]$ (2pyNO = *t*-butyl 2-pyridyl nitroxide) with a relatively high barrier of magnetization reversal,” R. Murakami, T. Ishida, S. Yoshii, and H. Nojiri, *Dalton Trans.*, **42**, 13968-13973 (2013).
- 133) “Synthesis, Structure, Luminescent and Magnetic Properties of Carbonato-Bridged $\text{Zn}^{\text{II}}_2\text{Ln}^{\text{III}}_2$ Complexes $[(\mu_4\text{-CO}_3)_2\{\text{Zn}^{\text{II}}\text{L}^1\text{Ln}^{\text{III}}(\text{NO}_3)_2\}_2]$ ($\text{Ln}^{\text{III}} = \text{Gd}^{\text{III}}, \text{Tb}^{\text{III}}, \text{Dy}^{\text{III}}$; $\text{L}^1 = \text{N,N}'\text{-Bis(3-methoxy-2-oxybenzylidene)-1,3-propanediaminato}$, $\text{L}^2 = \text{N,N}'\text{-Bis(3-ethoxy-2-oxybenzylidene)-1,3-propanediaminato}$),” K. Ehama, Y. Ohmichi, S. Sakamoto, T. Fujinami, N. Matsumoto, N. Mochida, T. Ishida, Y. Sunatsuki, M. Tsuchimoto, and N. Re, *Inorg. Chem.*, **52**, 12828-12841 (2013).
- 134) “Determination of Crystal-Field Energy Levels and Temperature Dependence of Magnetic Susceptibility for Dy^{3+} in $[\text{Dy}_2\text{Pd}]$ Heterometallic Complex,” M. Karbowski, C. Rudowicz, and T. Ishida, *Inorg. Chem.*, **52**, 13199-13206 (2013).
- 135) “Exchange Couplings and Its Chemical Trend Studied by High-Frequency EPR on Heterometallic $[\text{Ln}_2\text{Ni}]$ Complexes,” A. Okazawa, T. Shimada, N. Kojima, S. Yoshii, H. Nojiri, and T. Ishida, *Inorg. Chem.*, **52**, 13351-13355 (2013).
- 136) “Preparation and Characterization of $[\text{Gd}(\text{hfac})_3(\text{DTBN})(\text{H}_2\text{O})]$ (DTBN = Di-*t*-butyl Nitroxide). Ferromagnetic $\text{Gd}^{3+}\text{-Gd}^{3+}$ Super-superexchange,” T. Kanetomo and T. Ishida, *Chem. Commun.*, **50**, 2529-2531 (2014).
- 137) “Multifunctional π -Expanded Macrocyclic Oligothiophene 6-Mers and Related Macrocyclic Oligomers,” M. Iyoda, K. Tanaka, H. Shimizu, M. Hasegawa, T. Nishiuchi, S. Sugibayashi, T. Nishinaga, Y. Kunugi, T. Ishida, H. Otani, H. Sato, K. Inukai, K. Tahara, and Y. Tobe, *J. Am. Chem. Soc.*, **136**, 2389-2396 (2014).
- 138) “Doubly TEMPO-Coordinated Gadolinium(III), Lanthanum(III), and Yttrium(III) Complexes. Strong Superexchange Coupling across Rare Earth Ions,” R. Murakami, T. Nakamura, and T. Ishida, *Dalton Trans.*, **43**, 5893-5898 (2014).
- 139) “Synthesis, Structure, Luminescence, and Magnetic Properties of a Single Ion Magnet, ‘*mer*’-[Tris(*N*-[(imidazol-4-yl)-methylidene]-DL-phenylalaninato)terbium(III) and Related DL-Alaninato Derivative,” S. Yamauchi, T. Fujinami, N. Matsumoto, N. Mochida, T. Ishida, Y. Sunatsuki, M. Watanabe, M. Tsuchimoto, C. Coletti, and N. Re, *Inorg. Chem.*, **53**, 5961-5971 (2014).
- 140) “Linker Stoichiometry Controlled Stepwise Supramolecular Growth of a Flexible Cu_2Tb Single Molecule Magnet from Monomer to Dimer to 1D Chain,” S. Ghosh, Y. Ida, T. Ishida, and A. Ghosh, *Cryst. Growth Desn.*, **14**, 2588-2598 (2014).
- 141) “Metal-Organic Frameworks from Two-dimensional Copper(II) Complexes Bridged with *N*-Methylpyrazine-2-carboxamide,” M. Yoshida, T. Kanetomo, H. Muroyama, T. Ishida, and T. Kogane, *Polyhedron*, **81**, 493-498 (2014).

- 142) “Polymeric One- and Two-dimensional Copper(I) Iodide Complexes Showing Photoluminescence Tunable by Azaaromatic Ligands,” N. Kitada and T. Ishida, *CrystEngComm*, **16**, 8035-8040 (2014).
- 143) “Synthesis, Structure, Luminescent and Magnetic Properties of Single Ion Magnets of Terbium(III) and Dysprosium(III) Complexes with a Triimidazolyl Tripodal Ligand and an Acetate,” S. Shintoyo, K. Murakami, T. Fujinami, N. Matsumoto, N. Mochida, T. Ishida, Y. Sunatsuki, M. Watanabe, M. Tsuchimoto, J. Mrozinski, C. Coletti, and N. Re, *Inorg. Chem.*, **53**, 10359-10369 (2014).
- 144) “Strongest Exchange Coupling in Gadolinium(III) and Nitroxide Coordination Compounds.” T. Kanetomo and T. Ishida, *Inorg. Chem.*, **53**, 10794-10796 (2014).
- 145) “Synthesis of Mixed-Valence Hexanuclear Mn(II/III) Clusters from Its Mn(II) Precursor: Variations of Catecholase-like Activity and Magnetic Coupling,” P. Kar, Y. Ida, T. Kanetomo, M. G. B. Drew, T. Ishida, and A. Ghosh, *Dalton Trans.*, **44**, 9795-9804 (2015).
- 146) “Relationship Between Torsion and Anisotropic Exchange Coupling in a Tb(III)-Radical Complex,” M. L. Baker, T. Tanaka, R. Murakami, S. Ohira-Kawamura, K. Nakajima, T. Ishida, and H. Nojiri, *Inorg. Chem.*, **54**, 5732-5738 (2015).
- 147) “Intermediate-Paramagnetic Phases with a Half and a Quarter Spin Entities in Fluorinated Biphenyl-3,5-diyl Bis(*tert*-butyl Nitroxides),” T. Konno, H. Kudo, and T. Ishida, *J. Mater. Chem. C*, **3**, 7813-7818 (2015).
- 148) “First Iron(II) Spin-crossover Complex with an N₅S Coordination Sphere,” M. Yamasaki and T. Ishida, *Chem. Lett.*, **44**, 920-921 (2015).
- 149) “Heating-Rate Dependence of Spin-Crossover Hysteresis Observed in an Iron(II) Complex Having Tris(2-pyridyl)methanol,” M. Yamasaki and T. Ishida, *J. Mater. Chem. C*, **3**, 7784-7787 (2015).
- 150) “Single-Molecule Magnet Involving Strong Exchange Coupling in Terbium(III) Complex with 2, 2'-Bipyridin-6-yl *tert*-Butyl Nitroxide,” T. Kanetomo, S. Yoshii, H. Nojiri, and T. Ishida, *Inorg. Chem. Front.*, **2**, 860-866 (2015).
- 151) “Strong Ferromagnetic Exchange Interactions in Hinge-like Dy(O₂Cu)₂ Complexes Involving Double Oxygen Bridges,” Y. Ida, S. Ghosh, A. Ghosh, H. Nojiri, and T. Ishida, *Inorg. Chem.*, **54**, 9543-9555 (2015).
- 152) “Spin-Crossover Hysteresis of [Fe^{II}(L_H^{iPr})₂(NCS)₂] (L_H^{iPr} = *N*-2-Pyridylmethylene-4-Isopropylaniline) Accompanied by Isopropyl Conformation Isomerism,” N. Mochida, A. Kimura, and T. Ishida, *Magnetochemistry*, **1**, 17-27 (2015).
- 153) “A Designed Room-Temperature Triplet Ligand from Pyridine-2,6-diyl Bis(*tert*-butyl Nitroxide),” H. Kawakami, A. Tonegawa, and T. Ishida, *Dalton Trans.*, **45**, 1306-1309 (2016) (Journal Inside Front Cover).
- 154) “Trigonal Bipyramid Ni₃Cl₃O₂ Cluster Showing the High-Spin *S* = 3 Ground State with Uniaxial Magnetic Anisotropy,” Y. Ida, H. Nojiri, and T. Ishida, *Bull. Chem. Soc. Jpn.*, **89**, 2256-2266 (2016) (BCSJ Award; Open Access).
- 155) “Luminescent Dy^{III} Single Ion Magnets with Same N₆O₃ Donor Atoms but Different Donor Atom Arrangements, ‘*fac*’-[Dy^{III}(HL^{DL-ala})₃]·8H₂O and ‘*mer*’-[Dy^{III}(HL^{DL-phe})₃]·7H₂O,” D. Hamada, T. Fujinami, S. Yamaguchi, N. Matsumoto, N. Mochida, T. Ishida, Y. Sunatsuki, M. Tsuchimoto, C. Coletti, and N. Re, *Polyhedron*, **109**, 120-128 (2016).
- 156) “Ferromagnetic Chains of Ground Triplet 5-Methoxy-1,3-phenylene Bis(*tert*-butyl nitroxide),” T. Yoshitake and T. Ishida, *Chem. Lett.*, **45**, 391-393 (2016).
- 157) “Thermally Activated Paramagnets from Diamagnetic Polymers of Biphenyl-3,5-diyl Bis(*tert*-butyl Nitroxides) with Methyl and Fluoro Groups at the 2'- and 5'-Positions,” T. Yoshitake, H. Kudo, and T. Ishida, *Crystals*, **6**, 00030 (9 pages) (2016).
- 158) “Syntheses, Crystal Structure and Magnetic Properties of an Unprecedented One Dimensional Coordination Polymer Derived from {(NiL)₂Ln} Node and a Pyrazine Spacer (H₂L = *N,N'*-Bis(salicylidene)-1,3-propanediamine, Ln = Gd, Tb and Dy),” S. Ghosh, P. Mahapatra, T. Kanetomo, M. G. B. Drew, T. Ishida, and A. Ghosh, *ChemistrySelect*, **1**, 2722-2729 (2016).
- 159) “Strongest Ferromagnetic Coupling in Designed Gadolinium(III)-Nitroxide Coordination Compounds,” T. Kanetomo, T. Yoshitake, and T. Ishida, *Inorg. Chem.*, **55**, 8140-8146 (2016).

- 160) "Spectroscopic and Magnetic Studies of Lanthanoid-TEMPO Complexes as Potential Single-Molecule Magnets: Interplay of the Crystal-Field and Exchange Coupling Effects," M. Karbowiak, C. Rudowicz, T. Nakamura, R. Murakami, and T. Ishida, *Chem. Phys. Lett.*, **662**, 163-168 (2016).
- 161) "Iron(II) Complex Tripodally Chelated with 1,1,1-Tris(2-pyridyl)ethane Showing Room-Temperature Spin-Crossover," T. Ishida, T. Kanetomo, and M. Yamasaki, *Acta Crystallogr. C*, **72**, 797-801 (2016).
- 162) "Notably Strong Antiferromagnetic Interaction in a Methylene-Bridged Bis(dihydrophenanthridin-*N*-oxyl)," T. Kanetomo and T. Ishida, *Chem. Lett.*, **46**, 188-190 (2017).
- 163) "Circular and Chainlike Copper(II)-Lanthanide(III) Complexes Generated by Assembly Reactions of Racemic and Chiral Copper(II) Cross-Linking Ligand-Complex with $\text{Ln}^{\text{III}}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$ ($\text{Ln}^{\text{III}} = \text{Gd}^{\text{III}}, \text{Tb}^{\text{III}},$ and Dy^{III})," T. Ueno, T. Fujinami, N. Matsumoto, M. Furusawa, R. Irie, N. Re, T. Kanetomo, T. Ishida, Y. Sunatsuki, *Inorg. Chem.*, **56**, 1679-1695 (2017).
- 164) "A Ground Triplet Dinuclear Copper(II) Spiro-Complex Involving Tetraiminoneopentane," A. Kimura and T. Ishida, *Polyhedron*, **126**, 45-48 (2017).
- 165) "Giant Exchange Coupling Evidenced with a Magnetization Jump at 52 T for a Gadolinium-Nitroxide Chelate," T. Kanetomo, T. Kihara, A. Miyake, A. Matsuo, M. Tokunaga, K. Kindo, H. Nojiri, and T. Ishida, *Inorg. Chem.*, **56**, 3310-3314 (2017) (ACS Editors' Choice; Open Access).
- 166) "Forced Proximity of Nitroxide Groups in Pincer Compounds with a Xanthene Spacer," N. Koizumi and T. Ishida, *Tetrahedron Lett.*, in press (2017). doi: 10.1016/j.tetlet.2017.06.004
- 167) "Structural Variations in $(\text{CuL})_2\text{Ln}$ Complexes of a Series of Lanthanide Ions with Salen-Type Unsymmetrical Schiff Base (H_2L) and the Dy and Tb Derivatives as Potential Single-Molecule Magnets," P. Mahapatra, S. Ghosh, N. Koizumi, T. Kanetomo, T. Ishida, and A. Ghosh, submitted (2017).
- 168) "A New $S = 0 \rightleftharpoons S = 2$ "Spin-crossover" Scenario Found in a Nickel(II) Bis(nitroxide) System," Y. Homma and T. Ishida, submitted (2017).

B. 学会プロシーディングスなど

- 1) "Approaches from High-Spin Organic Molecules to Organic Ferromagnets," T. Ishida, K. Inoue, N. Koga, N. Nakamura, and H. Iwamura, *Mater. Res. Soc. Sym. Proc.*, **247**, 407-416 (1992).
- 2) "Syntheses, Second-Order NLO properties, and Crystal Structures of Tetrasubstituted Ethylene, Trisubstituted Schiff-Base, and Diphenoxymethane Derivatives," T. Ishida, H. Osaka, T. Nogami, R. Yamazaki, M. Yasui, F. Iwasaki, A. Mizoguchi, M. Kuhata, T. Uemiyama, and A. Nishimura, *Synth. Metals*, **56**, 2013-2018 (1993).
- 3) "Attempted Syntheses of Dibenzotetraoxafulvalene," T. Ishida, Y. Ishihara, T. Yamaga, and T. Nogami, *Synth. Metals*, **56**, 1968-1971 (1993).
- 4) "Synthesis of Multi-Sulfur TTF Derivatives and Conducting Organic Salts," T. Nogami, T. Ishida, H. Yoshikawa, M. Tanaka, T. Naito, H. Kobayashi, and A. Kobayashi, *Synth. Metals*, **56**, 2039-2043 (1993).
- 5) "Intermolecular Ferromagnetic Interaction of 4-(1-Pyrenylmethyleneamino)-2,2,6,6-tetramethylpiperidin-1-oxyl," T. Ishida, K. Tomioka, T. Nogami, H. Iwamura, K. Yamaguchi, W. Mori, and Y. Shirota, *Mol. Cryst. Liq. Cryst.*, **232**, 99-102 (1993).
- 6) "Intramolecular Magnetic Interaction of Transition Metal Ions in Complexes Containing Pyrimidine or Pyrazine as a Bridging Ligand," T. Ishida, S.-I. Mitsubori, T. Nogami, and H. Iwamura, *Mol. Cryst. Liq. Cryst.*, **233**, 345-350 (1993).
- 7) "Synthesis and Characterization of C_{60} Adducts with Benzyne, Dichlorocarbene, Dienes, and Oxygen," T. Nogami, M. Tsuda, T. Ishida, S. Kurono, and M. Ohashi, *Advanced Materials '93, I, B: Magnetic, Fullerene, Dielectric, Ferroelectric, Diamond and Related Materials, Trans. Mat. Res. Soc. Jpn.*, **Vol. 14B**, 1181-1184 (1994).
- 8) "New Organic Ferromagnets: 4-Arylmethyleneamino-2,2,6,6-tetramethylpiperidin-1-oxyl (Aryl = Phenyl, 4-Biphenyl, 4-Chlorophenyl, and 4-Phenoxyphenyl)," T. Nogami, K. Togashi, H. Tsuboi, T. Ishida, H. Yoshikawa, M. Yasui, F. Iwasaki, H. Iwamura, N. Takeda, and M. Ishikawa, *Synth. Metals*, **71**, 1813-1814 (1995).

- 9) "Intra- and Inter-Molecular Ferromagnetic Interaction of Transition Metal Complexes Containing Pyrimidine or Pyrazine Derivatives," T. Ishida, S.-i. Mitsubori, N. Takeda, M. Ishikawa, H. Iwamura, and T. Nogami, *Synth. Metals*, **71**, 1791-1792 (1995).
- 10) "Ferromagnetism of Pyrimidine-Bridged Copper(II) Complexes," T. Ishida, T. Nogami, M. Yasui, F. Iwasaki, H. Iwamura, N. Takeda, and M. Ishikawa, *Mol. Cryst. Liq. Cryst.*, **279**, 87-96 (1996).
- 11) "Ferromagnetism of Organic Radical Crystals of TEMPO Derivatives," T. Nogami, T. Ishida, M. Yasui, F. Iwasaki, H. Iwamura, N. Takeda, and M. Ishikawa, *Mol. Cryst. Liq. Cryst.*, **279**, 97-106 (1996).
- 12) "Crystal Structures of TEMPO Radicals Showing Ferromagnetic Interaction," M. Yasui, H. Yoshikawa, H. Yamamoto, T. Ishida, T. Nogami, and F. Iwasaki, *Mol. Cryst. Liq. Cryst.*, **279**, 77-85 (1996).
- 13) "Magnetic Properties of 4- and 5-Methylpyrimidine-Bridged Copper(II) Complexes," M. Nakagawa, Y. Ishikawa, T. Kogane, T. Ishida, M. Yasui, F. Iwasaki, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **286**, 29-34 (1996).
- 14) "Ferromagnetic Interaction in a Pyrimidine-Bridged Copper(II) Nitrate Complex," K. Nakayama, M. Nakagawa, T. Ishida, Y. Ishikawa, M. Yasui, F. Iwasaki, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **306**, 379-384 (1997).
- 15) "Ferromagnetism of Organic Radical Crystals of Perdeuterated TEMPO Derivatives," T. Nogami, R. Imachi, T. Ishida, N. Takeda, and M. Ishikawa, *Mol. Cryst. Liq. Cryst.*, **305**, 211-216 (1997).
- 16) "Magnetic Interaction via β -Hydrogen Atoms in TEMPO Derivatives," T. Kawakami, A. Oda, S. Takeda, W. Mori, T. Ishida, M. Yasui, F. Iwasaki, T. Nogami, and K. Yamaguchi, *Mol. Cryst. Liq. Cryst.*, **306**, 141-150 (1997).
- 17) "Solid State ^1H -MAS-NMR and Spin Densities on Protons of the Organic Ferromagnetic TEMPO Derivatives," G. Murata, S. Takeda, T. Kawakami, W. Mori, R. Imachi, T. Ishida, T. Nogami, and K. Yamaguchi, *Mol. Cryst. Liq. Cryst.*, **306**, 307-314 (1997).
- 18) "Ferromagnetism of Organic Radical Crystals of Deuterated TEMPO Derivatives," R. Imachi, T. Ishida, N. Takeda, M. Ishikawa, and T. Nogami, *Synth. Met.*, **85**, 1743-1744 (1997).
- 19) "Magnetic Properties of Pyrimidine-Bridged Copper(II) Complexes," T. Ishida, K. Nakayama, M. Nakagawa, W. Sato, Y. Ishikawa, M. Yasui, F. Iwasaki, and T. Nogami, *Synth. Met.*, **85**, 1655-1658 (1997).
- 20) "Electrical Conductivities of MOTT Salts," T. Nogami, K. Yoshida, T. Ishida, and A. Kobayashi, *Synth. Met.*, **86**, 2135-2136 (1997).
- 21) "Crystal Structures of 4-(4-Halobenzylideneamino)-TEMPO Radicals Showing Magnetic Interactions," F. Iwasaki, H. Yamamoto, J. H. Yoshikawa, K. Takada, E. Kan-nari, M. Yasui, T. Ishida, and T. Nogami, *Mol. Cryst. Liq. Cryst. Sci. Technol., Sect. A*, **313**, 223-228 (1998).
- 22) "Intermolecular Ferromagnetic Interaction in π -Conjugated Nitroxide Radical Crystals," T. Ishida, Y. Akui, S.-y. Masaoka, M. Nakagawa, D. Hashizume, M. Yasui, F. Iwasaki, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **334**, 89-98 (1999).
- 23) "Magnetic Phase Transition of Copper Complexes Containing dmit or TCNQ Anion," T. Ishida, Y. Kokubo, K. Nakayama, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **335**, 323-332 (1999).
- 24) "Single Crystal Magnetic Susceptibility of 1-D $[\text{PMCu}(\text{NO}_3)_2(\text{H}_2\text{O})_2]_n$," R. Feyerherm, T. Ishida, T. Nogami, and M. Steiner, *Mol. Cryst. Liq. Cryst.*, **335**, 235-244 (1999).
- 25) "Pressure Effects of Organic Ferromagnetic Radical $p\text{-Cl-C}_6\text{H}_4\text{-CH=N-TEMPO}$," M. Mito, T. Kawae, K. Takeda, T. Ishida, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **334**, 323-332 (1999).
- 26) "Spin Density of a Ferromagnetic Tempo Derivative," Y. Pontillon, T. Ishida, E. Lelíre-Berna, T. Nogami, E. Ressouche, and J. Schweizer, *Mol. Cryst. Liq. Cryst.*, **334**, 359-367 (1999).
- 27) "Possible Superhigh-Spin Cluster in the Crystals of a Novel Nitronyl Nitroxide Radical as Studied by FT Pulsed ESR," D. Shiomi, K. Sato, T. Takui, K. Itoh, M. Tamura, Y. Nishio, K. Kajita, M. Nakagawa, T. Ishida, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **335**, 359-369 (1999).
- 28) " μSR Studies on the Organic Radical Magnet, 4-Arylmethyleneamino-TEMPO," S. Ohira, T. Ishida, T. Nogami, I. Watanabe, F. L. Pratt, and K. Nagamine, *Physica B*, **289-290**, 123-127 (2000).
- 29) "Magnetism of Pyrimidine-Bridged Metal(II) Halide Complexes," K. Zusai, T. Kusaka, T. Ishida, R. Feyerherm, M. Steiner, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **343**, 127-132 (2000).

- 30) "Magnetic Phase Transition of Py_2N^- and NCNH^- -Bridged Nickel(II) Complexes at about 25 K," K. Zusai, T. Katayama, T. Ishida, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **343**, 121-126 (2000).
- 31) "Synthesis and Properties of New Oxygen-Containing Fulvalene Derivatives: Dibenzo- and Dinaphthotetraoxafulvalenes and Dibenzodioxadithiafulvalene," T. Nogami, K. Tanaka, T. Ishida, and A. Kobayashi, *Synth. Met.*, **120**, 755-756 (2001).
- 32) "Zero-Field Muon Spin Rotation Study on TEMPO-Based Magnets, Ar-CH=N-TEMPO," T. Ishida, S. Ohira, I. Watanabe, T. Nogami, and K. Nagamine, *Polyhedron*, **20**, 1545-1549 (2001).
- 33) "Ferromagnetic $S = 1$ Chain Formed by a Square Ni_2S_2 Motif in $\text{Ni}(\text{qt})_2$ (qt = Quinoline-8-thiolate). Magnetic Properties of Related Compounds," T. Miyake, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Polyhedron*, **20**, 1551-1555 (2001).
- 34) "Radical-Copper Hexanuclear Arrays Showing Ferromagnetic Interactions," J. Omata, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Polyhedron*, **20**, 1557-1561 (2001).
- 35) "Anomalous Magnetism in Organic Radical Ferromagnets, 4-Arylmethyleneamino-2,2,6,6-tetramethylpiperidin-1-yloxyl just above Tc Studied by μSR Method," S. Ohira, T. Ise, K. Nakayama, T. Ishida, T. Nogami, I. Watanabe, and K. Nagamine, *Polyhedron*, **20**, 1223-1227 (2001).
- 36) "Radical-Copper Macrocycles and Related Compounds," J. Omata, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **376**, 455-462 (2002).
- 37) "Magnetic Phase Transition of Fe^{II} , Co^{II} , and Ni^{II} Complexes Bridged by Pyrimidine and Dicyanamide," T. Kusaka, T. Ishida, D. Hashizume, F. Iwasaki, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **376**, 463-468 (2002).
- 38) "Solvated Magnets of $\text{Fe}[\text{N}(\text{CN})_2]_2$ (pyrimidine): Transition Phenomena Tuned by Guests," T. Kusaka, T. Ishida, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **379**, 259-264 (2002).
- 39) "First Isolation and Characterization of an HIN Complex, $\text{CdCl}_2(\text{HIN})_4$ (HIN = 4,4,5,5-Tetramethylimidazolin-1-oxyl)," T. Ise, T. Ishida, and T. Nogami, *Mol. Cryst. Liq. Cryst.*, **379**, 147-152 (2002).
- 40) "Magnetic Properties of 9,9-Disubstituted 9,10-Dihydroacridine-10-yloxyls," M. Seino, Y. Akui, T. Ishida, and T. Nogami, *Synth. Met.*, **133-134**, 581-583 (2003).
- 41) "Hybrid Magnetic Solids of Radical-Substituted Heteroaromatic Compounds with Metal Halides," T. Matsuyama, Y. Iwata, T. Ishida, and T. Nogami, *Synth. Met.*, **133-134**, 611-613 (2003).
- 42) "Low Temperature Deuterium NMR Studies on Magnetism of TEMPO Derivatives," G. Maruta, S. Takeda, M. Ooishi, R. Imachi, T. Ishida, and T. Nogami, *Synth. Met.*, **133-134**, 577-579 (2003).
- 43) "Hydroiminonitroxide Complexes with Metal Hexafluoroacetylacetonates," T. Ise, T. Ishida, and T. Nogami, *Synth. Met.*, **137**, 1281-1282 (2003).
- 44) "Host-Guest Chemistry of Radical-Copper Wheels. A Supramolecular Control of Magnetic Exchange Coupling," T. Ishida, J. Omata, and T. Nogami, *Polyhedron*, **22**, 2133-2138 (2003).
- 45) "*meta*-Phenylene-Bridged Bis(imino Nitroxide) Biradicals as Potential High-Spin Ligands," T. Ichimura, K. Doi, C. Mitsuhashi, T. Ishida, and T. Nogami, *Polyhedron*, **22**, 2557-2564 (2003).
- 46) "Effect of Staggered Field in $S = 1/2$ Antiferromagnetic Chain: Copper Pyrimidine," T. Asano, D. Nomura, Y. Inagaki, Y. Ajiro, H. Nojiri, Y. Narumi, K. Kindo, T. Ishida, and T. Nogami, *Physica B: Cond. Matter*, **329**, 1006-1007 (2003).
- 47) "High-Field Magnetization Process in the $S = 1/2$ Chain $[\text{PM}\cdot\text{Cu}(\text{NO}_3)_2\cdot(\text{H}_2\text{O})]_n$ (PM = pyrimidine)," A. Matsuo, Y. Narumi, T. Asano, Y. Ajiro, T. Ishida, T. Nogami, and K. Kindo, *J. Mag. Mag. Mater.*, **272-276**, 1081-1082 (2004).
- 48) "ESR of Quantum Spin Chains with Staggered Fields," T. Asano, Y. Ajiro, Y. Inagaki, H. Nojiri, T. Sakon, T. Ishida, T. Nogami, J. P. Boucher, and M. Motokawa, *J. Mag. Mag. Mater.*, **272-276**, e685-e686 (2004).
- 49) "Nano-Sized Tetralanthanoid-Copper Complex Ions Containing Oximate Bridges," S. Ueki, M. Sahlan, T. Ishida, and T. Nogami, *Synth. Met.*, **154**, 217-220 (2005).
- 50) "Structures and Magnetic Properties of Copper(II) Complexes Containing Tetrakis(2-pyridyl)methane," A. Okazawa, T. Ishida, and T. Nogami, *Polyhedron*, **24**, 2584-2587 (2005).

- 51) "Structure and Magnetic Properties of 3d-4f Heterometallic Complexes Containing Di-2-pyridyl Ketoximate. An Approach to Single-Molecule Magnets," F. Mori, T. Ishida, and T. Nogami, *Polyhedron*, **24**, 2588-2592 (2005).
- 52) "Magnetic Properties of Transition Metal Complexes with 2,2'-Bipyridin-5-yl *t*-Butyl Nitroxide," T. Shimada, T. Ishida, and T. Nogami, *Polyhedron*, **24**, 2593-2598 (2005).
- 53) "Effect of Staggered Field on the Magnetization Process of $S = 1/2$ Antiferromagnetic Chains," A. Matsuo, K. Hosokawa, Y. Narumi, K. Okunishi, N. Maeshima, T. Asano, Y. Ajiro, T. Ishida, T. Nogami, and K. Kindo, *Prog. Theor. Phys. Suppl.*, **159**, 158-162 (2005).
- 54) "ET and TTF Salts with Lanthanide Complex Ions Showing Frequency-Dependent ac Magnetic Susceptibility," S. Ueki, T. Ishida, T. Nogami, and M. Tamura, *Mol. Cryst. Liq. Cryst.*, **455**, 129-134 (2006).
- 55) "Mono- and Dinitroxide Radicals from 9,9'(10*H*,10'*H*)-Spirobiacridine: An Approach to a D_{2d} Triplet Biradical," T. Ishida, M. Ooishi, N. Ishii, H. Mori, and T. Nogami, *Polyhedron*, **26**, 1793-1799 (2007).
- 56) "Preparation, Structures, and Magnetic Properties of Precursors for Chelating Compounds of 2-Pyridyl Nitroxide Derivatives," A. Okazawa, T. Nogami, and T. Ishida, *Polyhedron*, **26**, 1965-1969 (2007).
- 57) "Tetranuclear Heterometallic Cycle Dy_2Cu_2 and the Corresponding Polymer Showing Slow Relaxation of Magnetization Reorientation," S. Ueki, A. Okazawa, T. Ishida, T. Nogami, and H. Nojiri, *Polyhedron*, **26**, 1970-1976 (2007).
- 58) "Structures and Magnetic Properties of One-dimensional Copper(II) Complexes Bridged with Diazaaromatic Rings," T. Kogane, N. Koyama, T. Ishida, and T. Nogami, *Polyhedron*, **26**, 1811-1819 (2007).
- 59) "Ferromagnetic Coupling of Copper(II) and Nickel(II) Complexes with a Cyclam-Based Paramagnetic Host," K. Igarashi, T. Nogami, and T. Ishida, *Polyhedron*, **28**, 1672-1677 (2009).
- 60) "Strong Intramolecular Ferromagnetic Couplings in Nickel(II) and Copper(II) Complexes Chelated with *tert*-Butyl 5-Methoxyphenyl-2-pyridyl Nitroxide," A. Okazawa, T. Nogami, and T. Ishida, *Polyhedron*, **28**, 1917-1921 (2009).
- 61) "Magnetic Properties and Exchange Couplings of One-Dimensionally Arrayed 4f-3d Heterometallic $[Ln_2Cu_2]_n$ Compounds," A. Okazawa, R. Watanabe, H. Nojiri, T. Nogami, and T. Ishida, *Polyhedron*, **28**, 1808-1813 (2009).
- 62) "Structures and Magnetic Properties of Imidazolate-Bridged Tetra- and Polynuclear Copper(II) Complexes," N. Koyama, R. Watanabe, T. Ishida, T. Nogami, and T. Kogane, *Polyhedron*, **28**, 2001-2009 (2009).
- 63) "2,2'-Bipyridine-6,6'-diyl Bisnitroxide as a Paramagnetic Host. Encapsulation of a Zinc(II) Ion," K. Koide and T. Ishida, *Polyhedron*, **30**, 3034-3037 (2011).
- 64) "Ferromagnetic Oxovanadium(IV) Complexes Chelated with Tetrahalosalen Ligands," K. Fujiwara and T. Ishida, *Polyhedron*, **30**, 3073-3078 (2011).
- 65) "Single-Molecule Magnet Behavior Enhanced by Magnetic Coupling between 4f-3d Spins," A. Okazawa, H. Nojiri, T. Ishida, and N. Kojima, *Polyhedron*, **30**, 3140-3144 (2011).
- 66) "Exchange Couplings in One-dimensionally Arrayed 4f-3d Heterometallic $[Ln_2Cu_2]_n$ Compounds. A Chemical Trend of the Coupling Parameter," A. Okazawa, K. Fujiwara, R. Watanabe, N. Kojima, S. Yoshii, H. Nojiri, and T. Ishida, *Polyhedron*, **30**, 3121-3126 (2011).
- 67) "Attempted Synthesis of Axial-Equatorial Pyrazine-Bridged Copper(II) Complexes toward Homometallic Ferromagnetic Compounds," R. Watanabe, T. Shimada, N. Koyama, T. Ishida, and T. Kogane, *Polyhedron*, **30**, 3165-3170 (2011).
- 68) "Magnetic Study on Radical-Gadolinium(III) Complexes. Relationship between the Exchange Coupling and Coordination Structure," T. Ishida, R. Murakami, T. Kanetomo, and H. Nojiri, *Polyhedron*, **66**, 183-187 (2013).
- 69) "Crystal Structures and Magnetic Properties of Two-Dimensional Copper(II) Complexes Bridged with Pyrazine-2-carboxamide," M. Yoshida, T. Shimada, T. Ishida, and T. Kogane, *Polyhedron*, **66**, 75-80 (2013).

- 70) "Spin-crossover Thermal Hysteresis and Light-Induced Effect on Iron(II) Complexes with Tripodal Tris(2-pyridyl)methanol," M. Yamasaki and T. Ishida, *Polyhedron*, **85**, 795-799 (2015).
- 71) "Strong Exchange Couplings in Lanthanide Complexes with an Aliphatic Nitroxide Radical 1,1,3,3-Tetramethylisoindolin-2-oxyl," T. Nakamura and T. Ishida, *Polyhedron*, **87**, 302-306 (2015).
- 72) "Structures and Magnetic Properties of Three Compounds from Imidazolate-Bridged Copper(II) Pyrazolylborate Complexes," T. Kogane, T. Kanetomo, H. Muroyama, and T. Ishida, *Polyhedron*, **89**, 149-154 (2015).
- 73) "Luminescent Single-Ion Magnets from Lanthanoid(III) complexes with Monodentate Ketone Ligand," T. Kanetomo and T. Ishida, *AIP Conf. Proc.*, **1709**, 020015 (6 pages) (2016).
- 74) "Magnetic Exchange Interaction in Gadolinium(III) Complex Having Aliphatic Nitroxide Radical TEMPO," T. Nakamura and T. Ishida, *AIP Conf. Proc.*, **1709**, 020016 (3 pages) (2016).
- 75) "Pyridine-2,6-diyl Dinitroxides as Room-Temperature Triplet Ligands," H. Kawakami, A. Tonegawa, and T. Ishida, *AIP Conf. Proc.*, **1709**, 020017 (6 pages) (2016).
- 76) "Development of Biradical Ligands toward [2x2] Molecular Grids," K. Ichihashi, Y. Mazaki, S. Ohba, and T. Ishida, *AIP Conf. Proc.*, **1807**, 020018 (7 pages) (2017).
- 77) "Structures and Magnetic Properties of Transition Metal Complexes Involving 2,2'-Bipyridin-6-yl Nitroxide," A. Ondo and T. Ishida, *AIP Conf. Proc.*, **1807**, 020023 (6 pages) (2017).
- 78) "Chemical Trend on the Lanthanide-Radical Exchange Coupling," T. Ishida, T. Nakamura, T. Kihara, and H. Nojiri, *Polyhedron*, in press (2017). doi: 10.1016/j.poly.2017.03.008
- 79) "Exchange-Coupled Terbium-Radical Complex Tb-phNO Showing Slow Reversal of Magnetization," T. Kanetomo and T. Ishida, *Polyhedron*, in press (2017). doi: 10.1016/j.poly.2017.02.031
- 80) "Study on Imidazolate-Bridged Divalent Copper Ion Complexes Capped with Tris(1-pyrazolyl)methane," T. Kogane, A. Ondo, M. Yamasaki, T. Kanetomo, and T. Ishida, *Polyhedron*, in press (2017). doi: 10.1016/j.poly.2017.03.045
- 81) "Spin-Parity Behavior in the Exchange-Coupled Lanthanoid-Nitroxide Molecular Magnets," T. Ishida, *IOP Conf. Ser.: Mater. Sci. Eng.*, **202**, 012001 (10 pages) (2017).

C. 総説及び著書

- 1) "Magnetic Properties and Nonlinear Optical Properties of Organic Materials," T. Nogami and T. Ishida, *New Functionality Materials, Vol. C, Synthetic Process and Control of Functionality Materials*, 383-388 (1993).
- 2) "Pyrimidine-Bridged Transition Metal Complexes as a Polymer Approach to Molecule-Based Ferromagnets," T. Ishida and T. Nogami, *Recent Res. Devel. Pure Appl. Chem.*, **1**, 1-15 (1997).
- 3) "Molecular Design and Synthesis for Organic Molecular Magnetism - Novel Ferromagnetic TEMPO Radical Crystals," T. Nogami and T. Ishida, "Molecular Magnetism - New Magnetic Materials," Ed. by K. Ito and M. Kinoshita, Kodansha-Gordon and Breach Science Publishers (2000), Chap. 5.1, pp. 207-215.
- 4) "超分子的に制御できる分子磁性 (Molecule-Based Magnets Tunable by Means of Supramolecular Chemistry)", 石田尚行・野上 隆, *日本応用磁気学会誌〜トピックス〜*, **27**, 811-816 (2003).
- 5) "Electron Donors from Natural Products: Donating and Ligating Abilities and Charge-Transfer Complex Formation of Imidazo[1,2-a]pyrazin-3(7H)-ones," M. Suzuki, T. Fujii, T. Nogami, T. Hirano, and T. Ishida, *Multifunctional Conducting Molecular Materials*, ed. by G. Saito, F. Wudl, R. C. Haddon, T. Katsumi, T. Enoki, H. E. Katz, and M. Maesato, pp. 55-58, RSC Publishing, U.K. (2007).
- 6) 「反応速度式の正しい求め方」 (講座「化学の大学入試問題を考えるための基本」), 石田尚行, *化学と教育*, **58**, 226-229 (2010).
- 7) 「希土類ナノ磁石の交換相互作用評価」 (フォーカス26 <第14回> : 成果事例クローズアップ (ナノテク融合技術支援センターによるイノベーション創出支援事業)), 石田尚行, 野尻浩之, 吉居俊輔, 田中豪, *Nanotech Japan Bull.* **2010**, 3; https://nanonet.nims.go.jp/ntjb_pdf/F26-14.pdf.
- 8) 「分子磁性研究の展開と深化: 4f-3d単分子磁石の周辺研究を一例として」、石田尚行, *電子スピンスサイエンス*, **10** (Autumn), 94-99 (2012) (**Journal Cover Picture**).

- 9) 「スピン化学が拓く分子磁性の新展開- 設計から機能化まで-」(CSJカレントレビュー第16号)、石田尚行、第17章 *f-d系およびf-p系単分子磁石*、pp. 166-171、化学同人(2014年8月)。
- 10) “Attraction of complex chemistry: Quest for innovative single-molecule magnets,” T. Ishida, “Topics” in UEC e-Bulletin, December 2016 issue. <http://www.ru.uec.ac.jp/e-bulletin/archives/2016/12/>.

D. 特許・知的財産所有権

- 1) “新規なフラーレンC₆₀誘導体及びその製法,” 川隅 正則・伊東 孝之・野上 隆・石田 尚行、特開平9-25246、特願平7-17883、三田工業株式会社、公開日：平成9年(1997)1月28日、出願日：平成7年(1995)7月14日。